

Radiopharmaceutical	Activity Above Which Instructions Are Required	Activity Above Which a Record is Required	Recommendations for Interruption of Breast-Feeding
I-131 NaI	0.0004 mCi	0.002 mCi	Complete cessation
I-123	0.5 mCi	3 mCi	
Tc-99m DTPA	30 mCi	150 mCi	
Tc-99m MAA	1.3 mCi	6.5 mCi	12.6 hours for 4 mCi
Tc-99m Pertechnetate	3 mCi	15 mCi	
Tc-99m DISIDA	30 mCi	150 mCi	
Tc-99m Sestamibi	30 mCi	150 mCi	
Tc-99m MDP	30 mCi	150 mCi	
Tc-99m PYP	25 mCi	120 mCi	
Tc-99m Red Blood Cells	10 mCi	50 mCi	6 hours for 20 mCi
Tc-99m Red Blood Cells	30 mCi	150 mCi	
Tc-99m Sulfur Colloid	7 mCi	35 mCi	6 hours for 12 mCi
Tc-99m DTPA Aerosol	30 mCi	150 mCi	
Tc-99m MAG3	30 mCi	150 mCi	
Tc-99m White Blood Cells	4 mCi	15 mCi	24 hours for 30 mCi, 12 hours for 12 mCi
Ga-67 Citrate	0.04 mCi	0.2 mCi	1 month for 4 mCi, 2 weeks for 1.3 mCi, 1 week for 0.2 mCi
In-111 White Blood Cells	0.2 mCi	1 mCi	1 week for 0.5 mCi
Tl-201 Chloride	1 mCi	5 mCi	2 weeks for 3 mCi

Reference: NRC NUREG-1556 Appendix U, October 2002